An algorithmic approach in constructing infinitely many even size graphs with local antimagic chromatic number 3

Dedicated to Prof. S. Arumugam on the occasion of his 80th birthday

Gee-Choon Lau

Abstract. An edge labeling of a connected graph G = (V, E) is said to be local antimagic if it is a bijection $f: E \to \{1, \ldots, |E|\}$ such that for any pair of adjacent vertices x and y, $f^+(x) \neq f^+(y)$, where the induced vertex label $f^+(x) = \sum f(e)$, with e ranging over all the edges incident to x. The local antimagic chromatic number of G, denoted by $\chi_{la}(G)$, is the minimum number of distinct induced vertex labels over all local antimagic labelings of G. In this paper, we first introduce an algorithmic approach to construct a family of infinitely many even size non-regular tripartite graphs with $t \ge 1$ component(s) in which every component is of odd order $p \ge 9$ and of size q = n(p+1) for $n \ge 2$. We show that every graph in this family has local antimagic chromatic number 3. We then allow the m-th component to have order $p_m \geq 9$ and size $n_m(p_m+1)$ for $n_m \geq 2, 1 \leq m \leq t$. We prove that every such graph with all components having same order and size also has local antimagic chromatic number 3. Lastly, we construct another family of infinitely many graphs such that different components may have different order and size all of which having local antimagic chromatic number 3. Consequently, many other families of (possibly disconnected) graphs with local antimagic chromatic number 3 are also constructed.

Key words and phrases: Local antimagic chromatic number, disconnected, non-regular, tripartite.

AMS (MOS) Subject Classifications: 05C78; 05C69.

References

- S. Arumugam, K. Premalatha, M. Bača and A. Semaničová-Feňovčíková, Local antimagic vertex coloring of a graph, *Graphs Com*bin., 33 (2017), 275–285.
- [2] M. Bača, A. Semaničová-Feňovčíková, and T.M. Wang, Local antimagic chromatic number for copies of graphs, *Mathematics*, 9, 1230 (2021), https://doi.org/10.3390/math9111230.
- [3] J. Bensmail, M. Senhaji and K. Szabo Lyngsie, On a combination of the 1-2-3 conjecture and the antimagic labelling conjecture, *Discrete Math. Theoret. Comput. Sci.*, **19**(1) (2017), Paper no. 21, 17 pp.
- [4] T.L Chan, G.C. Lau and W.C. Shiu, Complete solutions on local antimagic chromatic number of three families of disconnected graphs, *Commun. Comb. Optim.*, (2024), https://doi.org/10.22049/CCO .2024.29032.1818.
- [5] G.C. Lau, S.M. Lee, K. Schaffer and S.M. Tong, On k-step Hamiltonian bipartite and tripartite graphs, *Malaya J. Mat.*, 2(3) (2014), 180–187.
- [6] G.C. Lau, H.K. Ng and W.C. Shiu, Affirmative solutions on local antimagic chromatic number, *Graphs Combin.*, 36 (2020), 1337–1354.
- [7] G.C. Lau, W.C. Shiu and H.K. Ng, On local antimagic chromatic number of cycle-related join graphs, *Discuss. Math. Graph Theory*, 4(1) (2021), 133–152, https://doi.org/10.7151/dmgt.2177.
- [8] G.C. Lau and W.C. Shiu, On local antimagic chromatic number of lexicographic product graphs, Acta Math. Hungar., 169(1) (2023), 158–170.
- [9] J. Haslegrave, Proof of a local antimagic conjecture, (2017), https: //doi.org/10.23638/DMTCS-20-1-18.